



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

$$P_6(\mu) = \frac{1}{18}(231\mu^6 - 315\mu^4 + 105\mu^2 - 5).$$

$$P_4(\mu) = \frac{1}{8}(35\mu^4 - 30\mu^2 + 3). \quad P_2(\mu) = \frac{1}{2}(3\mu^2 - 1).$$

$$\begin{aligned} \therefore \sin^3 \theta \cos^3 \theta \sin^2 \phi \cos \phi &= -\frac{1}{4}(11\mu^3 - 3\mu)(1 - \mu^2)^{\frac{3}{2}} \cos 3\phi \\ &\quad - \frac{3}{4}\mu(1 - \mu^2)^{\frac{3}{2}} \cos 3\phi - \frac{1}{8}(33\mu^5 - 30\mu^3 + 5\mu)\sqrt{(1 - \mu^2)} \cos \phi \\ &\quad + \frac{1}{808}(7\mu^3 - 3\mu)\sqrt{(1 - \mu^2)} \cos \phi + \frac{1}{8}\mu\sqrt{(1 - \mu^2)} \cos \phi. \end{aligned}$$

PROBLEMS FOR SOLUTION.

ARITHMETIC.

162. Proposed by G. B. M. ZERR, A. M., Ph. D., Professor of Chemistry and Physics, The Temple College, Philadelphia, Pa.

A trolley road is built between two towns, and it is found that the gross annual receipts amount to 20% of the original cost; the annual cost of repairs is 2% of the original cost; and the working expenses is \$3000 in addition to 20% of the net receipts. After a year a second road is built at the same cost as the first and it is found that the gross receipts and working expenses per year are doubled, while the cost of repairs for the new road is 1% of cost. If the net receipts for both roads is \$72,500, find the cost of each road, and the net receipts the first year.

163. Proposed by CHRISTIAN HORNING, A.M., Professor of Mathematics, Heidelberg University, Tiffin, O.

Three Dutchmen and their wives went to market to buy hogs. The names of the men were Hans, Klaus and Hendricks, and of the women, Gertrude, Anna and Katrine; but it was not known which was the wife of each man. They each bought as many hogs as each man or woman paid shillings for each hog, and each man spent three guineas more than his wife. Hendricks bought 23 hogs more than Gertrude, and Klaus bought 11 more than Katrine. What was the name of each man's wife?

ALGEBRA.

166. Proposed by MARCUS BAKER, U. S. Geological Survey, Washington, D. C.

$$\begin{aligned} \text{Solve} \quad ax + by &= 2xz \dots [1]. \\ cy + dz &= 2xy \dots [2]. \\ ez + fx &= 2yz \dots [3]. \end{aligned}$$

167. Proposed by G. B. M. ZERR, A. M., Ph. D., Professor of Chemistry and Physics, The Temple College, Philadelphia, Pa.

A weight of m pounds falls and is broken into n pieces after which it is found that all weights, in pounds, from 1 to m can be weighed. Find the weight of each piece. Apply when $m=121$, $n=5$.